

Highly Active Antiretroviral Therapy Use and HIV Transmission Risk Behaviors Among Individuals Who Are HIV Infected and Were Recently Released From Jail

Kristen Clements-Nolle, PhD, MPH, Rani Marx, PhD, MPH, Michael Pendo, MPH, Eileen Loughran, BA, Milton Estes, MD, and Mitchell Katz, MD

We evaluated highly active antiretroviral therapy (HAART) use and risk behaviors among 177 inmates who were HIV infected and were released and reincarcerated in San Francisco, Calif, jails over a 12-month period. During the month preceding reincarceration, HIV transmission risk behaviors were common among respondents, and 59% of those with a history of antiretroviral use were not taking HAART. HAART discontinuation was independently associated with homelessness, marijuana use, injection drug use, and not receiving community medical care. Postrelease interventions for inmates who are HIV infected are needed. (*Am J Public Health*. 2008;98:661–666. doi:10.2105/AJPH.2007.112656)

A quarter of the adults who are HIV infected in the United States pass through correctional facilities annually.¹ Incarceration represents an opportunity to provide critical HIV prevention and health services, including highly active antiretroviral therapy (HAART).^{2–6} The structured environment of jail and prison promote high rates of medication adherence,^{7–9} leading to favorable virological and immunologic outcomes.^{8,10–12}

However, 2 recent studies documented rising viral loads and falling CD4 cell counts among prisoners who were released and then reincarcerated.^{11,12} These results suggest that offenders may have difficulty transitioning from a situation in which antiretroviral therapy is taken in a structured and often directly observed setting to one in which medications must be self-managed while facing the challenges of community reentry.¹² Homelessness, joblessness, drug use, mental illness, and inconsistent health care among recently released inmates who are HIV infected likely contribute to discontinuation of or low adherence to treatment.^{5,11,12}

To our knowledge, no studies to date have directly assessed HAART use and adherence or factors associated with HAART use among recently released inmates. Furthermore, few studies have described the postrelease sexual and drug-related behaviors in this population.^{13,14} To better understand antiretroviral use and HIV transmission behaviors of recently released inmates, we studied 177 inmates who were HIV infected and were released and then reincarcerated in the San Francisco, Calif, jail system in a 12-month period.

METHODS

Participants and Recruitment

The San Francisco Department of Public Health's Forensic AIDS Project provides medical care, case management, discharge planning, and community referrals for all inmates who are HIV infected and who were identified through voluntary testing and inmate disclosure. Inmates taking HAART are released with a 7-day medication supply and a 1-month refill prescription. In 2001, a demonstration grant^{15–17} enhanced Forensic AIDS Project services by linking inmates who are HIV infected with community case managers before release and providing 5 months of postrelease supportive services.^{16,17}

From April 2001 through June 2003, Forensic AIDS Project providers approached all inmates who were HIV infected to determine participation interest. Interested individuals were referred to research associates who confirmed eligibility, obtained written informed consent, and conducted interviews in private rooms. Inmates were eligible if they

(1) were at least 18 years old, (2) spoke English or Spanish, and (3) had at least 2 weeks of jail time remaining in their sentence. Those with a planned transfer to another correctional facility or a documented mental health condition requiring community psychiatric placement were ineligible.

Analyses were restricted to 177 inmates who had been released from San Francisco jails at any point during the 12 months prior to the first day of their current incarceration (68% of the study participants).

Measures

The interview assessed sociodemographics, incarceration history, and use of services the month preceding reincarceration. The date participants last visited a community provider for HIV-related medical care was compared with their release date to determine the proportion receiving postrelease care. Participants prioritized the service they thought they would need most when released.

The Center for Epidemiological Studies Depression Scale was used to screen for depression¹⁸ with the standard cutoff score of 16 or higher.¹⁹ Two or more positive responses on the CAGE Questionnaire indicated past or present alcohol dependence.^{20,21} Sexual and drug use behaviors the month preceding reincarceration were measured. Serodiscordant unprotected sexual intercourse included any anal or vaginal intercourse with partners of opposite or unknown HIV status when a condom was not used.

Participants were given a pill card with pictures to determine antiretroviral use during the month preceding reincarceration. Antiretroviral use was classified as HAART according to recommendations from the International AIDS Society–USA Panel.^{22,23} Participants were also asked how many times a week they missed at least 1 dose or did not take a full dose of medication (dichotomized as never or less than once a week vs once a week or more).

Statistical Methods

Among participants with a history of antiretroviral therapy use ($n=108$), we used χ^2 analyses to assess factors associated with lack of HAART use during the month preceding reincarceration. We entered variables

TABLE 1—Characteristics of 177 HIV-Positive Jail Inmates Who Were Released and Reincarcerated in a 12-Month Period: San Francisco, Calif, January 2000 to June 2003

	No. (%) or Median (Range)
Demographics	
Median age, y	39.5 (22–62)
Gender	
Men	145 (82)
Women	20 (11)
Transgender, male-to-female	12 (7)
Race/Ethnicity	
African American	95 (54)
White	58 (33)
Hispanic	18 (10)
Other ^a	6 (3)
Education ^b	
Less than high school or general equivalency diploma	62 (35)
High school or general equivalency diploma	63 (36)
More than high school	51 (29)
Homeless ^c (month preceding reincarceration)	94 (53)
Health insurance ^b	
None	81 (48)
Public	85 (50)
Private	4 (2)
ADAP or Medicaid coverage for HIV medications (month preceding reincarceration)	108 (61)
Median monthly income (month preceding reincarceration), ^{b,d} \$	900 (0–24 000)
Documented AIDS diagnosis	60 (34)
Median jail and prison incarcerations ^b	17 (3–130)
Community service use	
Receipt of HIV medical care ^b (between release and reincarceration)	106 (61)
Case management visit ^b (month preceding reincarceration)	68 (39)
Mental health provider visit (month preceding reincarceration)	31 (18)
Alcohol or drug treatment (month preceding reincarceration)	50 (28)
Anticipated postrelease service needs ^b	
Housing	97 (55)
HIV medical care	22 (13)
Substance abuse treatment	16 (9)
Case management	11 (6)
Other services ^e	29 (17)
Depression and alcohol drug dependence	
Depression: Center for Epidemiological Studies Depression Scale score $\geq 16^b$	137 (78)
Alcohol dependence: CAGE score ≥ 2	75 (42)
Drug use (month preceding reincarceration)	
Crack cocaine (noninjection)	103 (58)
Marijuana (nonprescription)	90 (51)
Methamphetamine (noninjection)	46 (26)
Heroin (noninjection)	22 (12)
Cocaine (noninjection)	14 (8)
Amyl nitrites	8 (5)
Club drugs (MDMA, GHB, ketamine, LSD)	6 (3)

Continued

associated with lack of HAART use in bivariate analyses ($P \leq .05$) into a logistic regression model simultaneously; age, gender, race/ethnicity, and AIDS diagnosis were forced into the model. A correlation matrix of all multivariate parameter estimates detected no serious multicollinearity; deviance and Pearson residuals confirmed good model fit. Analyses were conducted with SAS version 9.0 (SAS Institute Inc, Cary, NC).

RESULTS

Overall, participants were economically disadvantaged, and repeat incarceration was common. More than half of the participants were homeless the month preceding reincarceration, and housing was their most pressing postrelease service need. Community behavioral risk taking included serodiscordant unprotected sexual intercourse (27%–38%) and distributive syringe sharing (17%). Almost two thirds had a history of antiretroviral use ($n = 108$); however, 59% of this group ($n = 64$) did not use HAART the month preceding reincarceration. Among HAART users ($n = 44$), 52% missed medication doses once a week or more (Table 1).

In the multivariate model, lack of HAART use during the month preceding reincarceration was significantly associated with homelessness, marijuana use, injection drug use, and not receiving community medical care (Table 2).

DISCUSSION

Although nearly three fourths of the inmates who are HIV infected in San Francisco jails who need antiretroviral therapy take it,¹⁰ we observed alarmingly low rates of HAART use and adherence among inmates who were released into the community and then reincarcerated. Our results support results of recent studies that documented loss of clinical benefits from HAART postrelease.^{11,12} Suboptimal HAART use while in the community compromises the personal health of ex-offenders and may increase HIV transmission,^{24–26} including transmission of drug-resistant strains.^{27,28} This is particularly troubling given the rates of serodiscordant

TABLE 1—Continued

Injection drug use ^b	91 (52)
Receptive syringe sharing ^f	22 (24)
Distributive syringe sharing ^{b,f}	15 (17)
Sharing other drug equipment ^f	31 (34)
Sexual behaviors	
Men who had sex with men (month preceding reincarceration)	
Serodiscordant/unknown UAI ^g	11 (27)
Men who had sex with women (month preceding reincarceration)	
Serodiscordant/unknown UVI ^h	23 (28)
Women who had sex with men (month preceding reincarceration)	
Serodiscordant/unknown UVI ^h	13 (38)
Male-to-female transgender people who had sex with men (month preceding reincarceration)	
Serodiscordant/unknown UAI ^g	8 (38)
Antiretroviral therapy history	
History of antiretroviral therapy ^b	
Yes	108 (62)
No	65 (38)
HAART use (month preceding reincarceration) ⁱ	
Yes	44 (41)
No	64 (59)
Missed HAART doses (month preceding reincarceration) ^{b,j}	
Never or less than once a week	20 (48)
More than once a week	22 (52)

Note. ADAP = AIDS Drug Assistance Program; MDMA = methylenedioxymethamphetamine, GHB = γ -hydroxybutyrate, LSD = lysergic acid diethylamide; UAI = unprotected anal intercourse; UVI = unprotected vaginal intercourse; HAART = highly active antiretroviral therapy.

^a“Other” included American Indian (4), Asian/Pacific Islander (1), and Creole (1).

^bMissing data: education (1), health insurance (7), median income (4), median lifetime incarcerations (8), community HIV medical care (3), community case management (2), history of antiretroviral use (4), HAART adherence (2), depression (2), injection drug use (1), distributive syringe sharing (1).

^cHomeless was defined as having spent at least 1 night on streets, in parks, in abandoned buildings, or in shelters during the month preceding reincarceration.

^dCombines all illegal and legal sources of income, including benefits (before taxes).

^eOther services included food assistance, education, job training, financial benefits, mental health care, and family services; each represented less than 5% of the sample.

^fAmong injection drug users ($n = 91$).

^gAmong men who had sex with men ($n = 41$) and transgender persons who had sex with men ($n = 8$).

^hAmong men who had sex with women ($n = 81$) and women who had sex with men ($n = 13$).

ⁱAmong those who ever used antiretroviral therapy ($n = 108$).

^jAmong those who used HAART the month preceding reincarceration ($n = 44$).

unprotected sexual intercourse and distributive syringe sharing we observed and others have reported.^{14,29–31}

Consistent with studies among the urban poor, we identified several barriers to post-release HAART use that may be amenable to intervention, including homelessness,^{32,33} injection drug use,^{34–36} marijuana use,³⁷ and inconsistent medical care^{33,38,39}; however, we did not find an association with depression.^{34,40–42} Many ex-offenders have limited access to food, refrigeration, and a safe place to store

medications,⁴³ and the provision of stable housing may be a necessary first step⁴⁴ toward sustaining postrelease HAART use. Also, more drug treatment options^{11,45–47} are urgently needed, and comprehensive programs that link inmates who are HIV infected to health services,^{48–52} support medication adherence,⁴³ and reduce behavioral risk taking^{13,53–55} are essential.

Our study may have overestimated community risk, because the month preceding reincarceration is often unstable and chaotic.⁵⁴

Additionally, research has shown that use of self-report to document HIV medication adherence is limited,⁵⁶ and temporality could not be established with our cross-sectional design.

Despite these limitations, our results highlight the need for coordinated public health interventions that begin during incarceration and continue postrelease. Such interventions are critical to improving health outcomes for inmates who are HIV infected and preventing further HIV transmission in the community. ■

About the Authors

At the time of this study, the authors were with the San Francisco Department of Public Health, San Francisco, CA.

Requests for reprints should be sent to Kristen Clements-Nolle, PhD, MPH, University of Nevada, Reno, School of Public Health/274, Reno, NV 89557 (e-mail: clements@unr.edu).

This brief was accepted August 7, 2007.

Contributors

K. Clements-Nolle designed the research study, conducted the data analysis, and drafted the brief. R. Marx contributed to study design and assisted with writing and revising the brief. M. Pendo directed the study and assisted with writing early drafts of the brief. E. Loughran coordinated field aspects of the study, conducted chart reviews for analysis, and contributed to revisions of the brief. M. Estes coordinated participant recruitment and contributed to revisions of the brief. M. Katz provided guidance on study design and data analysis and assisted with writing and revising the brief.

Acknowledgments

The Corrections Demonstration Grant, jointly funded by the Centers for Disease Control and Prevention (CDC) and the Health Resources and Services Administration, funded the intervention. Supplemental evaluation funding was provided by the state of California, Office of AIDS, and the CDC.

We appreciate assistance in all aspects of the study from Continuum HIV Day Services; San Francisco Department of Public Health, Forensic AIDS Project; San Francisco Department of Public Health, AIDS Office; and the San Francisco Sheriff's Department. Specifically, we would like to thank Cicily Emerson for study coordination; Naomi Akers, Ari Bachrach, Octavia Baker, Alice Heimsoth, Jessie James, Jelousy Jiggets, Zappa Montag, and Jessie Wofsy for participant interviewing; and Hilda Erlenbach, Christine Geoghegan, Shoshana Hall, Susan Halperin, and Sharon Hofer for participant recruitment.

We would like to dedicate this brief to Michael Pendo (August 10, 1964–November 15, 2005), who directed the study and was committed to improving the health of inmates and other underserved populations.

TABLE 2—Factors Associated With Lack of Highly Active Antiretroviral Therapy (HAART) Use the Month Preceding Reincarceration in Jail: San Francisco, Calif, January 2000 to June 2003

	No HAART Use (n = 64), ^a No. (%)	HAART Use (n = 44), ^a No. (%)	AOR (95% CI) (n = 106) ^b
Age			
≤ Median (39.5 y)	34 (53)	19 (43)	0.59 (0.22, 1.53)
> Median	30 (47)	25 (57)	Reference
Gender			
Men	48 (75)	36 (82)	0.39 (0.12, 1.26)
Women or transgender	16 (25)	8 (18)	Reference
Race			
African American	31 (48)	27 (61)	1.15 (0.43, 3.09)
Other	33 (52)	17 (39)	Reference
Education			
Less than a high school or general equivalency diploma	28 (44)	13 (30)	...
High school or general equivalency diploma	36 (56)	30 (70)	
Homeless			
Yes	38 (59)	16 (36)*	2.71 (1.04, 7.08)*
No	26 (41)	28 (64)	Reference
Monthly income			
≤ Median (\$900)	31 (48)	26 (62)	...
> Median	33 (52)	16 (38)	
AIDS diagnosis			
Yes	32 (50)	21 (48)	1.78 (0.67, 4.75)
No	32 (50)	23 (52)	Reference
No. of times previously incarcerated			
≤ Median (17 times)	27 (43)	18 (45)	...
> Median	36 (57)	22 (55)	
Received postrelease HIV medical care			
Yes	36 (57)	35 (81)	Reference
No	27 (43)	8 (19)**	3.08 (1.05, 9.05)*
Case management visit			
Yes	26 (41)	16 (37)	...
No	37 (59)	27 (63)	
Mental health provider visit			
Yes	9 (14)	9 (20)	...
No	55 (86)	35 (80)	
Alcohol or drug treatment			
Yes	16 (25)	13 (30)	...
No	48 (75)	31 (70)	
Depression (Center for Epidemiological Studies Depression Scale score ≥ 16)			
Yes	51 (81)	34 (79)	...
No	12 (19)	9 (21)	

Continued

Human Participant Protection

All study protocols and materials received approval from the Committee on Human Research at the University of California, San Francisco, and the Data Governance Committee of San Francisco General Hospital. In addition, a Certificate of Confidentiality protecting study participants was obtained from the Centers for Disease Control and Prevention.

References

- Hammett TM, Harmon MP, Rhodes W. The burden of infectious disease among US correctional facilities, 1997. *Am J Public Health*. 2002;92:1789–1794.
- Glaser JB, Greifinger RB. Correctional health care: a public health opportunity. *Ann Intern Med*. 1993;118:139–145.
- Flanigan TP, Rich JD, Spaulding A. HIV among incarcerated persons: a missed opportunity. *AIDS*. 1999;13:2475–2476.
- Hammett TM, Gaiter JL, Crawford C. Reaching seriously at-risk populations: health interventions in criminal justice settings. *Health Educ Behav*. 1998;25:99–120.
- Spaulding A, Stephenson B, Macalino G, Ruby W, Clarke JG, Flanigan TP. Human immunodeficiency virus in correctional facilities. *Clin Infect Dis*. 2002;35:305–312.
- Boutwell A, Rich JD. HIV infection behind bars. *Clin Infect Dis*. 2004;38:1761–1763.
- Altice FL, Mostashari F, Friedland GH. Trust and acceptance of and adherence to antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2001;28:47–58.
- Kirkland LR, Fischl MA, Tashima KT, et al. Response to lamivudine-zidovudine plus Abacavir twice daily in antiretroviral-naïve, incarcerated patients with HIV infection taking directly observed treatment. *Clin Infect Dis*. 2002;34:511–518.
- Wohl DA, Stephenson BL, Golin CE, et al. Adherence to directly observed antiretroviral therapy among human immunodeficiency virus-infected prison inmates. *Clin Infect Dis*. 2003;36:1572–1576.
- White MC, Mehrotra A, Menendez E, Estes M, Goldenson J, Tulskey JP. Jail inmates and HIV care: provision of antiretroviral therapy and *Pneumocystis carinii* pneumonia prophylaxis. *Int J STD AIDS*. 2001;12:380–385.
- Springer SA, Pesanti E, Hodges J, Macura T, Doros G, Altice F. Effectiveness of antiretroviral therapy among HIV-infected prisoners: re-incarceration and the lack of sustained benefits and release to the community. *Clin Infect Dis*. 2004;38:1754–1760.
- Stephenson BL, Wohl DA, Golin CE, Tien HC, Stewart P, Kaplan AH. Effect of release from prison and re-incarceration on the viral loads of HIV-infected individuals. *Public Health Rep*. 2005;120:84–88.
- Grinstead O, Zack B, Faigues B. Reducing postrelease risk behavior among HIV seropositive

TABLE 3—Continued

Alcohol abuse (CAGE score ≥ 2) ^c			
Yes	26 (41)	12 (27)	...
No	38 (59)	32 (73)	
Marijuana use (nonprescription)			
Yes	42 (66)	15 (34)***	3.63 (1.39, 9.52)**
No	22 (34)	29 (66)	Reference
Crack cocaine use (noninjection)			
Yes	43 (67)	24 (55)	...
No	21 (33)	20 (45)	
Methamphetamine use (noninjection)			
Yes	19 (30)	6 (14)*	1.31 (0.40, 4.30)
No	45 (70)	38 (86)	Reference
Injection drug use			
Yes	41 (64)	19 (44)*	3.17 (1.16, 8.67)*
No	23 (36)	24 (56)	Reference

Note. AOR = adjusted odds ratio; CI = confidence interval. Ellipsis indicates that the variable was not entered in multivariate mode.

^aWe used the χ^2 test to assess group differences in HAART use among 108 participants with a history of any antiretroviral use; 4 participants had missing data for antiretroviral use.

^bBased on logistic regression with simultaneous entry (probability of not using HAART modeled); 2 participants were not included because of missing data on 1 or more variables.

^cCAGE is a 4-question alcoholism screening instrument whose name is a mnemonic designating an individual who has attempted to "cut down" on alcohol consumption, is "annoyed" by criticism of his or her drinking, feels "guilty," and needs an "eye-opener" drink in the morning. The presence of 2 or more of these characteristics is considered indicative of an alcohol use disorder.

* $P \leq .05$; ** $P \leq .01$; *** $P \leq .001$.

drug-naïve HIV-1-infected persons in 10 US cities. *J Infect Dis.* 2004;189:2174–2180.

29. Latka MH, Metsch LR, Mizuno Y, et al. Unprotected sex among HIV-positive injection drug-using women and their serodiscordant male partners. *J Acquir Immune Defic Syndr.* 2006;42:222–228.

30. Weinhardt LS, Kelly JA, Brondino MJ, et al. HIV transmission risk behavior among men and women living with HIV in 4 cities in the United States. *J Acquir Immune Defic Syndr.* 2004;36:1057–1066.

31. Kral AH, Lorvick J, Ciccarone D, et al. HIV prevalence and risk behaviors among men who have sex with men and inject drugs in San Francisco. *J Urban Health.* 2005;82:43–50.

32. Knowlton A, Arnsten J, Eldred L, et al. Individual, interpersonal, and structural correlates of effective HAART use among urban active injection drug users. *J Acquir Immune Defic Syndr.* 2006;41:486–492.

33. Pulvirenti JJ, Glowacki R, Muppiddi U, et al. Hospitalized HIV-infected patients in the HAART era: a view from the inner city. *AIDS Patient Care STDS.* 2003;17:565–573.

34. Moss AR, Hahn JA, Perry S, et al. Adherence to highly active antiretroviral therapy in the homeless population in San Francisco: a prospective study. *Clin Infect Dis.* 2004;39:1190–1198.

35. Howard AA, Arnsten JH, Lo Y, et al. A prospective study of adherence and viral load in a large multi-center cohort of HIV-infected women. *AIDS.* 2002;16:2175–2182.

36. Arnsten JH, Demas PA, Grant RW, et al. Impact of active drug use on antiretroviral therapy adherence and viral suppression in HIV-infected drug users. *J Gen Intern Med.* 2002;17:377–381.

37. Kalichman SC, Rompa D. HIV treatment adherence and unprotected sex practices in people receiving antiretroviral therapy. *Sex Transm Infect.* 2003;79:59–61.

38. Keruly JC, Conviser R, Moore RD. Association of medical insurance and other factors with receipt of antiretroviral therapy. *Am J Public Health.* 2002;92:852–857.

39. Celentano DD, Galai N, Sethi AK. Time to initiating highly active antiretroviral therapy among HIV-infected injection drug users. *AIDS.* 2001;15:1707–1715.

40. Wadrop-Valverde D, Valverde E. Homelessness and psychological distress as contributors to antiretroviral nonadherence in HIV-positive injecting drug users. *AIDS Patient Care STDS.* 2005;19:326–334.

41. Anastos K, Schneider MF, Gange SJ, et al. The association of race, sociodemographic, and behavioral characteristics with response to highly active antiretroviral therapy in women. *J Acquir Immune Defic Syndr.* 2005;39:537–544.

42. Cook JA, Cohen MH, Burke J, et al. Effects of depressive symptoms and mental health quality of life on use of highly active antiretroviral therapy among HIV-seropositive women. *J Acquir Immune Defic Syndr.* 2002;30:401–409.

prison inmates: the health promotion program. *AIDS Educ Prev.* 2001;13:109–119.

14. Stephenson BL, Wohl DA, McKaig R, et al. Sexual behavior of HIV-seropositive men and women following release from prison. *Int J STD AIDS.* 2006;17:103–108.

15. Rapposelli KK, Kennedy MG, Miles JR, et al. HIV/AIDS in correctional settings: a salient priority for the CDC and HRSA. *AIDS Educ Prev.* 2002;14:103–113.

16. Robillard AG, Garner JE, Laufer FN, et al. CDC/HRSA HIV/AIDS intervention, prevention, and continuity of care demonstration project for incarcerated individuals within correctional settings and the community: part I, a description of corrections demonstration project activities. *J Correctional Health Care.* 2003;9:453–486.

17. Laufer FN, Arriola KR, Dawson-Rose CS, Kumarevelu K, Rapposelli KK. From jail to community: innovative strategies to enhance continuity of HIV/AIDS care. *Prison J.* 2002;82:84–100.

18. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977;1:385–401.

19. Weissman MM, Sholomskas D, Pottenger M, Prusoff BA, Locke BZ. Assessing depressive symptoms in five psychiatric populations: a validation study. *Am J Epidemiol.* 1977;106:203–214.

20. Ewing JA. Detecting alcoholism: the CAGE questionnaire. *JAMA.* 1984;252:1905–1907.

21. Fiellin DA, Reid MC, O'Connor PG. Screening for alcohol problems in primary care: a systematic review. *Arch Intern Med.* 2000;160:1977–1989.

22. Carpenter CC, Cooper DA, Fischl MA, et al. Antiretroviral therapy in adults. *JAMA.* 2000;283:381–391.

23. Yeni PG, Hammer SM, Carpenter CC, et al. Antiretroviral treatment for adult HIV infection in 2002. *JAMA.* 2002;288:222–235.

24. Musicco A, Lazzarin A, Nicolosi A, et al. Antiretroviral treatment of men infected with human immunodeficiency virus type 1 reduces the incidence of heterosexual transmission. *Arch Intern Med.* 1994;154:1971–1976.

25. Porco TC, Martin JN, Page-Shafer KA, et al. Decline in HIV infectivity following the introduction of highly active antiretroviral therapy. *AIDS.* 2004;18:81–88.

26. Castilla J, Del Romero J, Hernando V, Marinovich B, Garcia S, Rodriguez C. Effectiveness of highly active antiretroviral therapy in reducing heterosexual transmission of HIV. *J Acquir Immune Defic Syndr.* 2005;40:96–101.

27. Grant RM, Hecht FM, Warmerdam M, et al. Time trends in primary HIV-1 drug resistance among recently infected persons. *JAMA.* 2002;288:181–188.

28. Weinstock HS, Zaidi I, Heneine W, et al. The epidemiology of antiretroviral drug resistance among

43. Bamberger JD, Unick J, Klein P, Fraser M, Chesney M, Katz MH. Helping the urban poor stay with antiretroviral HIV drug therapy. *Am J Public Health*. 2000;90:699–701.
44. Tsemberis S, Gulcur L, Nakae M. Housing first, consumer choice, and harm reduction for homeless individuals with a dual diagnosis. *Am J Public Health*. 2004;94:651–656.
45. National Center on Addiction and Substance Abuse at Columbia University (CASA). *Behind Bars: Substance Abuse and America's Prison Population*. New York, NY: CASA; 1998.
46. Smith-Rohrberg D, Bruce RD, Altice FL. Review of corrections-based therapy for opiate-dependent patients: implications for buprenorphine treatment among correctional populations. *J Drug Issues*. 2004;34:451–480.
47. Hammett TM. HIV/AIDS and other infectious diseases among correctional inmates: transmission, burden, and an appropriate response. *Am J Public Health*. 2006;96:974–978.
48. Rich JD, Holmes L, Salas C, et al. Successful linkage of medical care and community services for HIV-positive offenders being released from prison. *J Urban Health*. 2001;78:279–289.
49. Richie BE, Freudenberg N, Page J. Reintegrating women leaving jail into urban communities: a description of a model program. *J Urban Health*. 2001;78:290–303.
50. Conklin TJ, Lincoln T, Flanagan TP. A public health model to connect correctional health care with communities. *Am J Public Health*. 1998;88:1249–1250.
51. Kim JY, Rich J, Zeiler S, et al. Successful community follow-up and reduced recidivism of HIV positive women prisoners. *J Correctional Health Care*. 1997;4:1–9.
52. Flanagan TP, Kim JY, Zierler S, Rich JD, Vigilante K, Bury-Maynard DA. A prison release program for HIV-positive women: linking them to health services and community follow-up. *Am J Public Health*. 1996;86:886–887.
53. Comfort M, Grinstead OA, Faigles B, Zack B. Reducing HIV risk among women visiting their incarcerated male partners. *Crim Justice Behav*. 2000;27:57–71.
54. Grinstead OA, Faigles B, Comfort M, et al. HIV, STD, and hepatitis risk to primary female partners of men being released from prison. *Women Health*. 2005;41:63–80.
55. Myers J, Zack B, Kramer K, Gardner M, Rucobo G, Costa-Taylor S. Get Connected: an HIV prevention case management program for men and women leaving California prisons. *Am J Public Health*. 2005;95:1682–1684.
56. Bangsberg DR, Hecht FM, Charlebois ED, et al. Adherence to protease inhibitors, HIV-1 viral load, and development of drug resistance in an indigent population. *AIDS*. 2000;14:357–366.